

Preventing pressure ulcers in adults


NICE Pathways bring together all NICE guidance, quality standards and other NICE information on a specific topic.

NICE Pathways are interactive and designed to be used online. They are updated regularly as new NICE guidance is published. To view the latest version of this pathway see:

<http://pathways.nice.org.uk/pathways/pressure-ulcers>

Pathway last updated: 22 February 2016

This document contains a single pathway diagram and uses numbering to link the boxes to the associated recommendations.



1 Adult at risk of developing a pressure ulcer

No additional information

2 Patient and carer information

Offer timely, tailored information to people who have been assessed as being at high risk of developing a pressure ulcer, and their family or carers. The information should be delivered by a trained or experienced healthcare professional and include:

- the causes of a pressure ulcer
- the early signs of a pressure ulcer
- ways to prevent a pressure ulcer
- the implications of having a pressure ulcer (for example, for general health, treatment options and the risk of developing pressure ulcers in the future).

Demonstrate techniques and equipment used to prevent a pressure ulcer.

Take into account individual needs when supplying information to people with:

- degenerative conditions
- impaired mobility
- neurological impairment
- cognitive impairment
- impaired tissue perfusion (for example, caused by peripheral arterial disease).

Quality standards

The following quality statement is relevant to this part of the interactive flowchart.

7. Information on preventing pressure ulcers

3 Risk assessment

Carry out and document an assessment of pressure ulcer risk for adults

- being admitted to secondary care or care homes in which NHS care is provided

- or receiving NHS care in other settings (such as primary and community care and emergency departments) if they have a risk factor, for example:
 - significantly limited mobility (for example, people with a spinal cord injury)
 - significant loss of sensation
 - a previous or current pressure ulcer
 - nutritional deficiency
 - the inability to reposition themselves
 - significant cognitive impairment.

Consider using a validated scale to support clinical judgement (for example, the Braden scale, the Waterlow score or the Norton risk-assessment scale) when assessing pressure ulcer risk.

Reassess pressure ulcer risk if there is a change in clinical status (for example, after surgery, on worsening of an underlying condition or with a change in mobility).

Quality standards

The following quality statements are relevant to this part of the interactive flowchart.

1. Pressure ulcer risk assessment in hospitals and care homes with nursing
2. Pressure ulcer risk assessment by community nursing services
3. Pressure ulcer risk reassessment

4 Skin assessment

Offer adults who have been assessed as being at high risk of developing a pressure ulcer a skin assessment by a trained healthcare professional (see [healthcare professional training and education \[See page 7\]](#) in this path). The assessment should take into account any pain or discomfort reported by the patient and the skin should be checked for:

- skin integrity in areas of pressure
- colour changes or discoloration (healthcare professionals should be aware that non-blanchable erythema may present as colour changes or discolouration, particularly in darker skin tones or types)
- variations in heat, firmness and moisture (for example, because of incontinence, oedema, dry or inflamed skin).

Use finger palpation or diascopy to determine whether erythema or discolouration (identified by skin assessment) is blanchable.

Start appropriate preventative action in adults who have non-blanching erythema and consider repeating the skin assessment at least every 2 hours until resolved.

Quality standards

The following quality statement is relevant to this part of the interactive flowchart.

4. Skin assessment

5 Care planning

Develop and document an individualised care plan for those who have been assessed as being at high risk of developing a pressure ulcer, taking into account:

- the outcome of risk and skin assessment
- the need for additional pressure relief at specific at-risk sites
- their mobility and ability to reposition themselves
- other comorbidities
- patient preference.

6 Barrier creams

Consider using a barrier preparation to prevent skin damage in adults who are at high risk of developing a moisture lesion or incontinence-associated dermatitis, as identified by skin assessment (such as those with incontinence, oedema, dry or inflamed skin).

7 Friction reduction

Medical technologies

NICE has published medical technologies guidance on [Parafricta Bootees and Undergarments to reduce skin breakdown in people with or at risk of pressure ulcers](#).

Resources

The following implementation tool is relevant to this part of the pathway.

[Parafricta Bootees and Undergarments to reduce skin breakdown in people with or at risk of pressure ulcers: costing statement](#)

8 Pressure redistribution

Use a high-specification foam mattress for adults who are:

- admitted to secondary care
- assessed as being at high risk of developing a pressure ulcer in primary and community care settings.

Consider a high-specification foam theatre mattress or an equivalent pressure redistributing surface for all adults who are undergoing surgery.

Consider the seating needs of people at risk of developing a pressure ulcer who are sitting for prolonged periods.

Consider a high-specification foam or equivalent pressure redistributing cushion for adults who use a wheelchair or who sit for prolonged periods.

Discuss with adults at high risk of developing a heel pressure ulcer and, where appropriate, their family and carers, a strategy to offload heel pressure, as part of their individualised care plan.

Quality standards

The following quality statement is relevant to this part of the interactive flowchart.

8. Pressure redistribution devices

9 Repositioning

Encourage adults who have been assessed as being at risk of developing a pressure ulcer to change their position frequently and at least every 6 hours. If they are unable to reposition

themselves, offer help to do so, using appropriate equipment if needed. Document the frequency of repositioning required.

Encourage adults who have been assessed as being at high risk of developing a pressure ulcer to change their position frequently and at least every 4 hours. If they are unable to reposition themselves, offer help to do so, using appropriate equipment if needed. Document the frequency of repositioning required.

Quality standards

The following quality statements are relevant to this part of the interactive flowchart.

5. Advice on repositioning
6. Help with repositioning

10 Interventions that should not be used

Skin massage

Do not offer skin massage or rubbing to prevent a pressure ulcer.

Nutritional supplements and hydration

Do not offer nutritional supplements specifically to prevent a pressure ulcer in adults whose nutritional intake is adequate.

Do not offer subcutaneous or intravenous fluids specifically to prevent a pressure ulcer in adults whose hydration status is adequate.

11 Healthcare professional training and education

Provide training to healthcare professionals on preventing a pressure ulcer, including:

- who is most likely to be at risk of developing a pressure ulcer
- how to identify pressure damage
- what steps to take to prevent new or further pressure damage
- who to contact for further information and for further action.

Provide further training to healthcare professionals who have contact with anyone who has been assessed as being at high risk of developing a pressure ulcer. Training should include:

- how to carry out a risk and skin assessment
- how to reposition
- information on pressure redistributing devices
- discussion of pressure ulcer prevention with patients and their carers
- details of sources of advice and support.

12 Managing pressure ulcers

[See Pressure ulcers / Managing pressure ulcers in adults](#)

Sources

Pressure ulcers (2014) NICE guideline CG179

Parafricta Bootees and Undergarments to reduce skin breakdown in people with or at risk of pressure ulcers (2014) NICE medical technologies guidance 20

Your responsibility

The guidance in this pathway represents the view of NICE, which was arrived at after careful consideration of the evidence available. Those working in the NHS, local authorities, the wider public, voluntary and community sectors and the private sector should take it into account when carrying out their professional, managerial or voluntary duties. Implementation of this guidance is the responsibility of local commissioners and/or providers. Commissioners and providers are reminded that it is their responsibility to implement the guidance, in their local context, in light of their duties to avoid unlawful discrimination and to have regard to promoting equality of opportunity. Nothing in this guidance should be interpreted in a way which would be inconsistent with compliance with those duties.

Copyright

Copyright © National Institute for Health and Care Excellence 2016. All rights reserved. NICE copyright material can be downloaded for private research and study, and may be reproduced for educational and not-for-profit purposes. No reproduction by or for commercial organisations, or for commercial purposes, is allowed without the written permission of NICE.

Contact NICE

National Institute for Health and Care Excellence
Level 1A, City Tower
Piccadilly Plaza
Manchester
M1 4BT

www.nice.org.uk

nice@nice.org.uk

0845 003 7781